



MARKSCHEME

May 1999

CHEMISTRY

Higher Level

Paper 3

OPTION C – HUMAN BIOCHEMISTRY


- C1. (a) Coconut oil [1 mark]
 The more unsaturated the oil, the bigger the iodine index (or converse). [1 mark]
- (b) M_R of oleic acid: $[(18 \times 12.0) + 34.0 + (16.0 \times 2)] = 282.0$ [1 mark]
 M_R of iodine: $126.9 \times 2 = 253.8$ [1 mark]
- Iodine Index = $\frac{100.0 \times 253.8}{282.0} = 90$ [1 mark]
- (c) Olive oil [1 mark]
highest degree of unsaturation/**most** unsaturated/**highest** iodine index [1 mark]
- (d) In order to **maintain body temperature/insulation** [1 mark]
 Energy source : or to protect organs [1 mark]
 Cell membranes to carry fat soluble vitamins [1 mark]
 to prevent skin drying/waterproofing
- Total [10 marks]**
- C2. (a) V_{max} ; maximum reaction rate. [1 mark]
- (b) 0–X: free active sites can accommodate increase in [subs] [1 mark]
 Thereafter: enzyme molecules are saturated with substrate/all active sites
 in use (so, they cannot act faster). [1 mark]
- (c) K_m represents [subs] at which the reaction rate is $\frac{1}{2}V_{max}$. [1 mark]
 The higher the value of K_m , the lower the activity of the enzyme.
 OR
 The lower the value of K_m , the higher the activity of the enzyme
 OR
 $K_m \propto \frac{1}{\text{activity}}$ of enzyme any one of these three answers [1 mark]
- (d) $\sim 3 \mu\text{mol dm}^{-3}$ [1 mark]
 accept between 2 and 4 (must give units for mark)
- Total [6 marks]**

- C3. (a) CH_2O or $(\text{CH}_2\text{O})_n$
 Carbonyl/ $\text{C}=\text{O}$ also aldehyde (alkanal)
 Hydroxyl/ OH


[3 marks]

(b)

OR




(D-glucose)




(L-glucose)

OR



(D-glucose)



(L-glucose)

must show 2nd* or 3rd OH is in a different direction from the other OH groups.



[1 mark]

In one isomer, the OH group on the C_1 carbon is in a different direction from that in the other isomer or shown by two correctly drawn ring structures.

[1 mark]

Optical isomerism/anomerism/stereoisomerism.

[1 mark]



(i.e. 1 mark for $\text{C}_{12}\text{H}_{22}\text{O}_{11}$ and 1 mark for balancing)

[2 marks]

Many monomers / monosaccharide molecules are involved

[1 mark]

OR Many C-O-C bonds formed

Total [9 marks]

OPTION D – ENVIRONMENTAL CHEMISTRY

- D1. (a)** $O_2 + (h\nu) \rightarrow 2O$ [1 mark]
- $O + O_2 \rightarrow O_3$ [1 mark]
- $O_3 \rightarrow O_2 + O$
or $O + O_3 \rightarrow 2O_2$ } do **not** give mark for $2O_3 \rightarrow 3O_2$ [1 mark]
- shorter** λ for (more) ozone formation (or converse) [1 mark]
- shorter** λ is higher energy (or converse) [1 mark]
- (b)** $CCl_2F_2 \rightarrow CClF_2 \cdot + Cl \cdot$ [1 mark]
- $O_3 + Cl \cdot \rightarrow ClO \cdot + O_2$ [1 mark]
- $ClO \cdot + O \rightarrow O_2 + Cl \cdot$ [1 mark]
- (c)** Catalysis on ice particles (surface). [1 mark]
- Total [9 marks]**
- D2. (a)** $CO_2(aq) + H_2O \rightleftharpoons H_2CO_3(aq)$ [1 mark], it forms an acid ([1 mark])
OR
 $CO_2 + H_2O \rightleftharpoons H^+(aq) + HCO_3^-(aq)$ [2 marks] as H^+ implies acidic } [2 marks]
- (b)** HNO_3 and H_2SO_4 / H_2SO_3 [2 marks]
[1] [1]
- NO – automobiles (some reaction between nitrogen and air) [1 mark]
(Subsequent reaction of the NO with oxygen to produce NO_2 and reaction of NO_2 with water.)
- SO_2 – smelters in the production of copper/power plants burning coal or oil (containing sulphur)/volcanoes [1 mark]
- equation for production of acid (any of the three acids) [1 mark]
- (c)** Leaching minerals from the soil
Fishless lakes
Damaging stone buildings
Damaging trees/forests
Iron/steel objects rust more quickly
Poorer health
- any two [1 mark] (accept valid alternatives) [1 mark]
- Total [8 marks]**

D3. Give [1 mark] for process and [1 mark] for what is removed

Primary: filtration
 sedimentation mostly insoluble material
 flocculation

any one method **[2 marks]**

Secondary: activated (bacterial) sludge oxidisable waste (organic products) **[2 marks]**

Tertiary: reverse osmosis
 electrodialysis soluble materials (NO₃⁻, detergents)
 chemical precipitation
 ion exchange

any one method **[2 marks]**

Due to greater levels of:
NO₃⁻/fertilisers
PO₄³⁻/detergents

any one by name, formula or source **[1 mark]**

reverse osmosis: semipermeable membrane and high pressure

electrodialysis: electrodes/cells and semipermeable membranes

chemical precipitation: chemical added combines with dissolved ions to give a precipitate (*e.g.* urea for nitrate)

ion exchange: unwanted ion is exchanged for a harmless ion

Also accept algae ponds and carbon beds.

any one **[1 mark]**

Total [8 marks]

OPTION E – CHEMICAL INDUSTRIES

- E1.** (a) (i) (Alumina has a very high melting point. By dissolving the alumina in cryolite) the electrolysis can be carried out at a much lower temperature/lowers melting point of alumina. [1 mark]
- (ii) $2\text{O}^{2-} - 4\text{e}^- \rightarrow \text{O}_2$ or the 'half' version $\text{O}^{2-} - 2\text{e}^- \rightarrow \frac{1}{2}\text{O}_2$ [1 mark]
- (iii) Oxygen produced at the anode reacts with the carbon electrode / Carbon (anode) burns in O_2 formed. *Must give $\text{C} + \text{O}_2 \rightarrow \text{CO}_2$ for the mark.* [1 mark]
- (b) (Alumina) reacts/dissolves with/in NaOH/KOH [1 mark]
 impurities/oxides do not react/dissolve [1 mark]
 Basic (must be included for the mark)
- (c) Examples could include:
 resistance to corrosion – window frames
 electrical conductivity **and** low density – overhead power cables
 low density – aircraft (fuselage) N.B. link between property and use essential
- accept valid alternatives, e.g. good heat conductor – saucepans (cooking) any two, [1 mark] each* [2 marks]
- (d) lower cost in energy terms/expensive to produce
 conserve resources
- accept other valid reasons* [2 marks]

Total [9 marks]

- E2.** (a) $(\text{RCOO})_2 \rightarrow (2\text{RCOO}\cdot) \rightarrow 2\text{R}\cdot + 2\text{CO}_2$ [1 mark]
 $\text{R}\cdot + \text{C}_2\text{H}_4 \rightarrow \text{RCH}_2\text{CH}_2\cdot$ [1 mark]
 $\text{RCH}_2\text{CH}_2\cdot + \text{C}_2\text{H}_4 \rightarrow \text{RCH}_2\text{CH}_2\text{CH}_2\text{CH}_2\cdot$ [1 mark]
 $\text{RCH}_2\text{CH}_2\text{CH}_2\text{CH}_2\cdot + \text{R}\cdot \rightarrow \text{R}(\text{CH}_2)_4\text{R}$ [1 mark]

If just states "initiation, propagation, termination", award [1 mark]

- Catalyst [1 mark]
 Ziegler [1 mark]
 Lower pressure/lower temperature [1 mark] [3 marks]


- (b) Not free radical/addition [1 mark]
 Monomers must contain two functional groups/small molecule (e.g. water) eliminated. [1 mark] [2 marks]

Total [9 marks]


- E3. (a)** Named fuel *e.g.* **propane** *[1 mark]*
Accept any named hydrocarbon
- balanced equation (*e.g.* $C_3H_8 + 5O_2 \rightarrow 3CO_2 + 4H_2O$) *[1 mark]*
- (b)** Balanced equation showing production of appropriate alkane and alkene
(*e.g.* $C_{10}H_{22} \rightarrow C_8H_{18} + C_2H_4$) *[2 marks]*
if only give alkane/alkene, [1 mark]
- (c)** Stage 1: **Fractional distillation** *[1 mark]*
Stage 2: Ethene is produced by **cracking** less valuable fractions *[1 mark]*
Stage 3: **Polymerisation** is used to convert ethene into polythene *[1 mark]*

Total [7 marks]

OPTION F – FUELS AND ENERGY



OR



not NO₂(g)

[1 mark]

(b) CO ensure efficient combustion (by monitoring combustion conditions and products)/ensuring excess air/oxygen

SO₂ (scrubbing the gas) by passing through an alkali/absorb in (slurry) powdered limestone – water/desulphurised coal OR fluid bed combustion.

not just scrubbing

NO₂ *if given in B* – (scrubbing the gas) by passing through an alkali

not just scrubbing


any one, [1 mark]

[1 mark]

(c) can be pumped
easy to burn/easy to mix with air/easier to control rate of combustion

any two

[2 marks]



[1 mark]

(ii) ΔH_f^0 (from Data Booklet) $-239 \text{ (kJ mol}^{-1}\text{)}$

[1 mark]

(Using $\text{CH}_3\text{OH} + 1\frac{1}{2}\text{O}_2 \rightarrow \text{CO}_2 + \text{H}_2\text{O}$)

$$\Delta H = H_{\text{products}}^0 - H_{\text{reactants}}^0$$

[1 mark]

$$= -(1 \times 393.5) - (2 \times 258.8) + (1 \times 239)$$

[1 mark]

$= -672.1 \text{ (kJ mol}^{-1}\text{)}$ *accept -672 kJ mol^{-1} (units must be included). If value for 2 moles given (-1344.2) give maximum of [3 marks].*

[1 mark]

correct answer scores [4 marks]

(N.B. last 3 marks could be obtained even with incorrect Data Booklet use *i.e.* last 3 marks consequential on **their** ΔH_f^0 value)

(iii) lower/less negative/more positive

[1 mark]

Energy needed to vapourise/evaporate the water/steam not condensed, therefore less energy released

[1 mark]

Total [11 marks]

F2. (a) Chemical (*which may be implicit*) rearrangement of outer electrons/no new elements or atoms formed.
(Nuclear – again may be implicit, *i.e.* order in question) change in nucleus emphasis/new elements/atoms formed/(outer) electron arrangement unchanged **[1 mark]**

- (b) shielding – to prevent escape of nuclear particles/radioactivity
control rods – to control and maintain a safe level of fission/control number of free neutrons
cooling system – maintain temperature of reactor (core)
moderator – to slow the neutrons

[2 marks] each **[6 marks]**

In the **absence** of **any** role being assigned give *2 marks* for three components (*1 mark* for two components)

Total [7 marks]

F3. (a) (i) (fast) electron
Sulphur **[2 marks]**

(ii) 5 half-lives
 5×14
= 70 days **[3 marks]**

- (b) – Surround wastes with concrete or other suitable material
– Solidify wastes, encapsulating them in glass or ceramic, then bury.
– Bury the wastes in an underground hole so that the wastes eventually melt and fuse with surrounding rock into a glassy ball
– Encase wastes in well-designed containers and drop them into the ocean
– Change harmful isotopes into harmless ones by using high-level neutron bombardment, lasers, or nuclear fusion


any two of the above, OR valid alternatives, [1 mark] each. **[2 marks]**

(N.B. searching for the principles)


Total [7 marks]

OPTION G – MODERN ANALYTICAL CHEMISTRY

G1. (a)



A



B

[2 marks]

(correct structural formulae but **incorrect** labelling deduct 1 mark).
In absence of 'labelling' assume order is as in question.

- (b) (i) Peaks in (intensity) ratio of **3:3:2**.
The chemical shifts of these hydrogens will be near **3.8 ppm**.
Any correct reference to splitting pattern, e.g. (smallest area peak) split into quartet/one peak (with greatest area) split into a triplet [3 marks]
- (ii) Peaks in (intensity) ratio of **6:1:1**.
The peak corresponding to the **most Hs will be split into a doublet**/one of the smallest area peaks will be a singlet/the other a septet (*accept complicated pattern*).
The O–H hydrogen will normally have a chemical shift at **4.5 ppm** [3 marks]
- (c) (i) No mark for just yes or no only. If YES, give [1 mark] for pointing out spectra are different. Award [3 marks] for a reasonable argument. If NO, give [1 mark] for pointing out spectra contain many similar peaks, e.g. 15, 29, 59. Only give further marks for thorough justification. [3 marks]
- (ii) [1 mark] for B (alcohol) having a higher boiling point. Give 2nd mark for hydrogen bonding in alcohols. [2 marks]

Total [13 marks]

- G2 (a) Vibrations (or stretching/bending) [1 mark]
 Change in dipole moment is required [1 mark]
 Different functional groups in different regions [1 mark]
 Precise absorption affected by neighbouring atoms (or mention of fingerprint region). [1 mark]
- (b) (i) ethanoic acid:
 C=O 1680–1750
 O–H 2500–3300/3580–3650
 C–H 2840–3095
- Any two correct for both marks* [2 marks]
- (ii) methyl methanoate:
 C=O 1680–1750
 C–H 2840–3095
 C–O 1000–1300 *any two, [1 mark] each* [2 marks]
- (c) O–H in ethanoic acid could be used/also C–O peak in ester could be used [1 mark]
 Other peaks/absorptions occur in both spectra [1 mark]
- (d) O–H [1 mark]
- $\frac{1}{3300} \text{ cm}^{-1} = 3.03 \times 10^{-4} \text{ cm}$ [1 mark]

If 'free' OH is not in the list, maximum energy will occur for C–H




$$\frac{1}{2480} \text{ cm}^{-1} = 4.03 \times 10^{-4} \text{ cm} \text{ (2 marks then possible)}$$

If free OH is in the list 1 mark possible for λ C–H = $4.03 \times 10^{-4} \text{ cm}$, i.e. wrong choice in (c) but converted correctly to λ value

Total [12 marks]



OPTION H – FURTHER ORGANIC CHEMISTRY

H1. (a) carbonyl, C=O (both needed) [1 mark] (accept alkanol/alkanone)




All three: [2 marks]. One/two: [1 mark]. Penalise extra structures once.


[3 marks]



[3 marks]




chiral C yet racemic mixture or words to that effect [1 mark]




continued...

Question H1. continued

N.B. Throughout (b)(i) and (b)(ii) only interested in  part of molecule.


Whatever else may be attached to the carbonyl carbon may be ignored.

Total [8 marks]



[1 mark]


[1 mark]




[1 mark]

[1 mark]


Either



[1 mark]




[1 mark]



[1 mark]

OR





[1 mark]

[2 marks] i.e. structure of ion [1 mark]
attack by Br^- [1 mark]

[3 marks]

Total [7 marks]

H3. (a) Bromination: Br_2 and AlBr_3 or FeBr_3 /reflux. **Must** include halogen carrier for mark.



[1 mark]

[2 marks]

* correct electrophile Br^+ [1 mark]; correct balanced equation [1 mark]

Nitration: Concentrated acids/heat




[1 mark]

[2 marks]

* correct electrophile NO_2^+ [1 mark]; correct balanced equation [1 mark]

(b)




[1 mark]

[1 mark]

[2 marks]

(c)



(Award [1 mark] for structure, [1 mark] for explanation in terms of stability of intermediate(s) etc. [$-\text{NO}_2$ is 3-diverting on its own no marks]).

[1 mark]

comment on directing influence (e.g. first NO_2 group e-withdrawing and deactivating, withdrawing e-density from 2 and 4 positions, so NO_2 group enters position 3). Remember some comment with some explanation expected, even NO_2 is 'meta' directing.

[1 mark]

Total [10 marks]