



International Baccalaureate®
Baccalauréat International
Bachillerato Internacional

Mathematics

Standard level

Specimen papers 1 and 2

For first examinations in 2014

CONTENTS

Mathematics standard level paper 1 specimen question paper

Mathematics standard level paper 1 specimen markscheme

Mathematics standard level paper 2 specimen question paper

Mathematics standard level paper 2 specimen markscheme



MATHEMATICS
STANDARD LEVEL
PAPER 1

SPECIMEN

1 hour 30 minutes

Candidate session number

0	0								
---	---	--	--	--	--	--	--	--	--

Examination code

X	X	X	X	-	X	X	X	X
---	---	---	---	---	---	---	---	---

INSTRUCTIONS TO CANDIDATES

- Write your session number in the boxes above.
- Do not open this examination paper until instructed to do so.
- You are not permitted access to any calculator for this paper.
- Section A: answer all questions in the boxes provided.
- Section B: answer all questions on the answer sheets provided. Write your session number on each answer sheet, and attach them to this examination paper and your cover sheet using the tag provided.
- At the end of the examination, indicate the number of sheets used in the appropriate box on your cover sheet.
- Unless otherwise stated in the question, all numerical answers should be given exactly or correct to three significant figures.
- A clean copy of the **Mathematics SL formula booklet** is required for this paper.
- The maximum mark for this examination paper is [90 marks].



0112


Full marks are not necessarily awarded for a correct answer with no working. Answers must be supported by working and/or explanations. Where an answer is incorrect, some marks may be given for a correct method, provided this is shown by written working. You are therefore advised to show all working.

SECTION A (44 Marks)

Answer **all** questions in the boxes provided. Working may be continued below the lines if necessary.

1. [Maximum mark: 7]

In the following diagram, $\mathbf{u} = \vec{AB}$ and $\mathbf{v} = \vec{BD}$.



The midpoint of \vec{AD} is E and $\frac{BD}{DC} = \frac{1}{3}$.

Express each of the following vectors in terms of \mathbf{u} and \mathbf{v} .

(a) \vec{AE} [3 marks]

(b) \vec{EC} [4 marks]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....




2. [Maximum mark: 5]

There are nine books on a shelf. For each book, x is the number of pages, and y is the selling price in pounds (£). Let r be the correlation coefficient.

(a) Write down the possible minimum and maximum values of r . [2 marks]

(b) Given that $r = 0.95$, which of the following diagrams best represents the data. [1 mark]



(c) For the data in diagram D, which **two** of the following expressions describe the correlation between x and y ?

- perfect, zero, linear, strong positive, strong negative,
 - weak positive, weak negative
- [2 marks]

.....

.....

.....

.....

.....



Do **NOT** write solutions on this page.

SECTION B (46 Marks)

Answer **all** the questions on the answer sheets provided. Please start each question on a new page.

8. [Maximum mark: 15]

Let $f(x) = 3(x+1)^2 - 12$.

(a) Show that $f(x) = 3x^2 + 6x - 9$. [2 marks]

(b) For the graph of f

(i) write down the coordinates of the vertex;

(ii) write down the y -intercept;

(iii) find both x -intercepts. [7 marks]

(c) **Hence** sketch the graph of f . [3 marks]

(d) Let $g(x) = x^2$. The graph of f may be obtained from the graph of g by the following two transformations

a stretch of scale factor t in the y -direction,

followed by a translation of $\begin{pmatrix} p \\ q \end{pmatrix}$.

Write down $\begin{pmatrix} p \\ q \end{pmatrix}$ and the value of t . [3 marks]



Do **NOT** write solutions on this page.

9. [Maximum mark: 14]

Two standard six-sided dice are tossed. A diagram representing the sample space is shown below.

		score on second die					
		1	2	3	4	5	6
score on first die	1	•	•	•	•	•	•
	2	•	•	•	•	•	•
	3	•	•	•	•	•	•
	4	•	•	•	•	•	•
	5	•	•	•	•	•	•
	6	•	•	•	•	•	•

Let X be the sum of the scores on the two dice.

(a) (i) Find $P(X = 6)$.

(ii) Find $P(X > 6)$.

(iii) Find $P(X = 7 | X > 6)$.

[6 marks]

(b) Elena plays a game where she tosses two dice.

If the sum is 6, she **wins** 3 points.

If the sum is greater than 6, she **wins** 1 point.

If the sum is less than 6, she **loses** k points.

Find the value of k for which the game is fair.


[8 marks]



Do **NOT** write solutions on this page.

10. [Maximum mark: 17]

Let $f(x) = \cos x + \sqrt{3} \sin x$, $0 \leq x \leq 2\pi$. The following diagram shows the graph of f .



The y -intercept is at $(0, 1)$, there is a minimum point at $A (p, q)$ and a maximum point at B .

(a) Find $f'(x)$. [2 marks]

(b) Hence

(i) show that $q = -2$;

(ii) verify that A is a minimum point. [10 marks]

(c) Find the maximum value of $f(x)$. [3 marks]

The function $f(x)$ can be written in the form $r \cos(x - a)$.

(d) Write down the value of r and of a . [2 marks]



Please **do not** write on this page.

Answers written on this page
will not be marked.





MARKSCHEME

SPECIMEN

MATHEMATICS

Standard Level

Paper 1

Instructions to Examiners

Abbreviations

- M** Marks awarded for attempting to use a correct **Method**.
- A** Marks awarded for an **Answer** or for **Accuracy**; often dependent on preceding **M** marks.
- R** Marks awarded for clear **Reasoning**.
- N** Marks awarded for **correct** answers if **no** working shown.
- AG** Answer given in the question and so no marks are awarded.

Using the markscheme

1 General

*Award marks using the annotations as noted in the markscheme eg **M1**, **A2**.*

2 Method and Answer/Accuracy marks

- Do **not** automatically award full marks for a correct answer; all working **must** be checked, and marks awarded according to the markscheme.
- It is generally not possible to award **M0** followed by **A1**, as **A** mark(s) depend on the preceding **M** mark(s), if any. An exception to this rule is when work for **M1** is missing, as opposed to incorrect (see point 4).
- Where **M** and **A** marks are noted on the same line, e.g. **M1A1**, this usually means **M1** for an **attempt** to use an appropriate method (e.g. substitution into a formula) and **A1** for using the **correct** values.
- Where there are two or more **A** marks on the same line, they may be awarded independently; so if the first value is incorrect, but the next two are correct, award **A0A1A1**.
- Where the markscheme specifies **M2**, **N3**, etc., do **not** split the marks, unless there is a note.
- Once a correct answer to a question or part-question is seen, ignore further working.

3 N marks

*If **no** working shown, award **N** marks for **correct** answers. In this case, ignore mark breakdown (**M**, **A**, **R**).*

- Do **not** award a mixture of **N** and other marks.
- There may be fewer **N** marks available than the total of **M**, **A** and **R** marks; this is deliberate as it penalizes candidates for not following the instruction to show their working.
- If a candidate has incorrect working, which somehow results in a correct answer, do **not** award the **N** marks for this correct answer. However, if the candidate has indicated (usually by crossing out) that the working is to be ignored, award the **N** marks for the correct answer.

4 Follow through marks (only applied after an error is made)

*Follow through (FT) marks are awarded where an incorrect answer from one **part** of a question is used correctly in **subsequent** part(s) or subpart(s). Usually, to award **FT** marks, **there must be working present** and not just a final answer based on an incorrect answer to a previous part. However, if the only marks awarded in a subpart are for the answer (i.e. there is no working expected), then **FT** marks should be awarded if appropriate.*

- Within a question part, once an **error** is made, no further **A** marks can be awarded for work which uses the error, but **M** marks may be awarded if appropriate.
- If the question becomes much simpler because of an error then use discretion to award fewer **FT** marks.
- If the error leads to an inappropriate value (e.g. probability greater than 1, use of $r > 1$ for the sum of an infinite GP, $\sin \theta = 1.5$, non integer value where integer required), do not award the mark(s) for the final answer(s).
- The markscheme may use the word “their” in a description, to indicate that candidates may be using an incorrect value.
- Exceptions to this rule will be explicitly noted on the markscheme.
- If a candidate makes an error in one part, but gets the correct answer(s) to subsequent part(s), award marks as appropriate, unless the question says hence. It is often possible to use a different approach in subsequent parts that does not depend on the answer to previous parts.

5 Mis-read

*If a candidate incorrectly copies information from the question, this is a mis-read (**MR**). Apply a **MR** penalty of 1 mark to that question*

- If the question becomes much simpler because of the **MR**, then use discretion to award fewer marks.
- If the **MR** leads to an inappropriate value (e.g. probability greater than 1, use of $r > 1$ for the sum of an infinite GP, $\sin \theta = 1.5$, non integer value where integer required), do not award the mark(s) for the final answer(s).
- Miscopying of candidates’ own work does **not** constitute a misread, it is an error.

6 Alternative methods

Candidates will sometimes use methods other than those in the markscheme. Unless the question specifies a method, other correct methods should be marked in line with the markscheme

- Alternative methods for complete questions are indicated by **METHOD 1, METHOD 2, etc.**
- Alternative solutions for part-questions are indicated by **EITHER . . . OR.**

7 Alternative forms

*Unless the question specifies otherwise, **accept** equivalent forms.*

- As this is an international examination, accept all alternative forms of **notation**.
- In the markscheme, equivalent **numerical** and **algebraic** forms will generally be written in brackets immediately following the answer.
- In the markscheme, **simplified** answers, (which candidates often do not write in examinations), will generally appear in brackets. Marks should be awarded for either the form preceding the bracket or the form in brackets (if it is seen).

8 Accuracy of Answers

If the level of accuracy is specified in the question, a mark will be allocated for giving the answer to the required accuracy.

There are 2 types of accuracy errors, and the final answer mark should not be awarded if these errors occur.

- **Rounding errors**: only applies to final answers not to intermediate steps.
- **Level of accuracy**: when this is not specified in the question the general rule applies to final answers: *unless otherwise stated in the question all numerical answers must be given exactly or correct to three significant figures.*

9 Calculators

No calculator is allowed. The use of any calculator on paper 1 is malpractice, and will result in no grade awarded.

10 Style

The markscheme aims to present answers using good communication, e.g. if the question asks to find the value of k , the markscheme will say $k = 3$, but the marks will be for the correct value 3 – there is usually no need for the “ $k =$ ”. In these cases, it is also usually acceptable to have another variable, as long as there is no ambiguity in the question, e.g. if the question asks to find the value of p and of q , then the student answer needs to be clear. Generally, the only situation where the full answer is required is in a question which asks for equations – in this case the markscheme will say “must be an equation”.

*The markscheme often uses words to describe what the marks are for, followed by examples, using the e.g. notation. These examples are not exhaustive, and examiners should check what candidates have written, to see if they satisfy the description. Where these marks are **M** marks, the examples may include ones using poor notation, to indicate what is acceptable.*

SECTION A

1. (a) $\vec{AE} = \frac{1}{2}\vec{AD}$ *A1*
attempt to find \vec{AD} *M1*
e.g. $\vec{AB} + \vec{BD}$, $\mathbf{u} + \mathbf{v}$
$$\vec{AE} = \frac{1}{2}(\mathbf{u} + \mathbf{v}) \left(= \frac{1}{2}\mathbf{u} + \frac{1}{2}\mathbf{v} \right)$$
 A1 *N2*
[3 marks]
- (b) $\vec{ED} = \vec{AE} = \frac{1}{2}(\mathbf{u} + \mathbf{v})$ *A1*
 $\vec{DC} = 3\mathbf{v}$ *A1*
attempt to find \vec{EC} *M1*
e.g. $\vec{ED} + \vec{DC}$, $\frac{1}{2}(\mathbf{u} + \mathbf{v}) + 3\mathbf{v}$
$$\vec{EC} = \frac{1}{2}\mathbf{u} + \frac{7}{2}\mathbf{v} \left(= \frac{1}{2}(\mathbf{u} + 7\mathbf{v}) \right)$$
 A1 *N2*
[4 marks]
Total [7 marks]
2. (a) min value of r is -1 , max value of r is 1 *A1A1* *N2*
[2 marks]
- (b) C *A1* *N1*
[1 mark]
- (c) linear, strong negative *A1A1* *N2*
[2 marks]
Total [5 marks]

3. (a) $4 \text{ (ms}^{-1}\text{)}$ *A1* *N1*
[1 mark]
- (b) recognising that acceleration is the gradient *M1*
e.g. $a(1.5) = \frac{4-0}{2-0}$
 $a = 2 \text{ (ms}^{-2}\text{)}$ *A1* *N1*
[2 marks]
- (c) recognizing area under curve *M1*
e.g. trapezium, triangles, integration
 correct substitution *A1*
e.g. $\frac{1}{2}(3+6)4, \int_0^6 |v(t)| dt$
 distance = 18 (m) *A1* *N2*
[3 marks]
- Total [6 marks]*
4. (a) (i) new mean is $20 + 10 = 30$ *A1* *N1*
- (ii) new sd is 6 *A1* *N1*
[2 marks]
- (b) (i) new mean is $20 \times 10 = 200$ *A1* *N1*
- (ii) **METHOD 1**
 variance is 36 *A1*
 new variance is $36 \times 100 = 3600$ *A1* *N2*
- METHOD 2**
 new sd is 60 *A1*
 new variance is $60^2 = 3600$ *A1* *N2*
[3 marks]
- Total [5 marks]*

5.	(a) attempt to use substitution or inspection	<i>M1</i>
	<i>e.g.</i> $u = 1 + e^x$ so $\frac{du}{dx} = e^x$	
	correct working	<i>A1</i>
	<i>e.g.</i> $\int \frac{du}{u} = \ln u$	
	$\ln(1 + e^x) + C$	<i>A1</i> <i>N3</i>
		<i>[3 marks]</i>
	(b) METHOD 1	
	attempt to use substitution or inspection	<i>M1</i>
	<i>e.g.</i> let $u = \sin 3x$	
	$\frac{du}{dx} = 3 \cos 3x$	<i>A1</i>
	$\frac{1}{3} \int u \, du = \frac{1}{3} \times \frac{u^2}{2} + C$	<i>A1</i>
	$\int \sin 3x \cos 3x \, dx = \frac{\sin^2 3x}{6} + C$	<i>A1</i> <i>N2</i>
		<i>[4 marks]</i>
	METHOD 2	
	attempt to use substitution or inspection	<i>M1</i>
	<i>e.g.</i> let $u = \cos 3x$	
	$\frac{du}{dx} = -3 \sin 3x$	<i>A1</i>
	$-\frac{1}{3} \int u \, du = -\frac{1}{3} \times \frac{u^2}{2} + C$	<i>A1</i>
	$\int \sin 3x \cos 3x \, dx = \frac{\cos^2 3x}{6} + C$	<i>A1</i> <i>N2</i>
		<i>[4 marks]</i>
	METHOD 3	
	recognizing double angle	<i>M1</i>
	correct working	<i>A1</i>
	<i>e.g.</i> $\frac{1}{2} \sin 6x$	
	$\int \sin 6x \, dx = \frac{-\cos 6x}{6} + C$	<i>A1</i>
	$\int \frac{1}{2} \sin 6x \, dx = -\frac{\cos 6x}{12} + C$	<i>A1</i> <i>N2</i>
		<i>[4 marks]</i>
		<i>Total [7 marks]</i>

6. (a) recognizing double angle *M1*
e.g. $3 \times 2 \sin x \cos x, 3 \sin 2x$
- $a = 3, b = 2$ *A1A1* *N3*
[3 marks]
- (b) substitution $3 \sin 2x = \frac{3}{2}$ *M1*
- $\sin 2x = \frac{1}{2}$ *A1*
- finding the angle *A1*
- e.g.* $\frac{\pi}{6}, 2x = \frac{5\pi}{6}$
- $x = \frac{5\pi}{12}$ *A1* *N2*

Note: Award *A0* if other values are included.

[4 marks]

Total [7 marks]

7. (a) $f'(x) = -x^{-2}$ (or $-\frac{1}{x^2}$) *A1* *N1*
- $f''(x) = 2x^{-3}$ (or $\frac{2}{x^3}$) *A1* *N1*
- $f'''(x) = -6x^{-4}$ (or $-\frac{6}{x^4}$) *A1* *N1*
- $f^{(4)}(x) = 24x^{-5}$ (or $\frac{24}{x^5}$) *A1* *N1*

[4 marks]

- (b) $f^{(n)}(x) = \frac{(-1)^n n!}{x^{n+1}}$ or $(-1)^n n! (x^{-(n+1)})$ *A1A1A1* *N3*


[3 marks]

Total [7 marks]

SECTION B

8. (a) $f(x) = 3(x^2 + 2x + 1) - 12$ *AI*
 $= 3x^2 + 6x + 3 - 12$ *AI*
 $= 3x^2 + 6x - 9$ *AG* *N0*
[2 marks]
- (b) (i) vertex is $(-1, -12)$ *A1A1* *N2*
- (ii) $y = -9$, or $(0, -9)$ *AI* *N1*
- (iii) evidence of solving $f(x) = 0$ *M1*
e.g. factorizing, formula
- correct working *AI*
e.g. $3(x + 3)(x - 1) = 0, x = \frac{-6 \pm \sqrt{36 + 108}}{6}$
- $x = -3, x = 1$, or $(-3, 0), (1, 0)$ *A1A1* *N2*
[7 marks]

(c)



A1A1A1 *N3*

Note: Award *AI* for a parabola opening upward,
AI for vertex in approximately correct position,
AI for intercepts in approximately correct positions.
 Scale and labelling not required.

[3 marks]

- (d) $\begin{pmatrix} p \\ q \end{pmatrix} = \begin{pmatrix} -1 \\ -12 \end{pmatrix}, t = 3$ *A1A1A1* *N3*
- [3 marks]*

Total [15 marks]

9. (a) (i) number of ways of getting $X = 6$ is 5 *A1*
 $P(X = 6) = \frac{5}{36}$ *A1* *N2*
- (ii) number of ways of getting $X > 6$ is 21 *A1*
 $P(X > 6) = \frac{21}{36} \left(= \frac{7}{12} \right)$ *A1* *N2*
- (iii) $P(X = 7 | X > 6) = \frac{6}{21} \left(= \frac{2}{7} \right)$ *A2* *N2*

[6 marks]

(b) attempt to find $P(X < 6)$ *M1*

e.g. $1 - \frac{5}{36} - \frac{21}{36}$

$P(X < 6) = \frac{10}{36}$ *A1*

fair game if $E(W) = 0$ (may be seen anywhere) *R1*

attempt to substitute into $E(X)$ formula *M1*

e.g. $3\left(\frac{5}{36}\right) + 1\left(\frac{21}{36}\right) - k\left(\frac{10}{36}\right)$

correct substitution into $E(W) = 0$ *A1*

e.g. $3\left(\frac{5}{36}\right) + 1\left(\frac{21}{36}\right) - k\left(\frac{10}{36}\right) = 0$

work towards solving *M1*

e.g. $15 + 21 - 10k = 0$

$36 = 10k$ *A1*

$k = \frac{36}{10} (= 3.6)$ *A1* *N4*

[8 marks]

Total [14 marks]

10. (a) $f'(x) = -\sin x + \sqrt{3} \cos x$ *A1A1* *N2*
[2 marks]
- (b) (i) at A, $f'(x) = 0$ *R1*
- correct working *A1*
 e.g. $\sin x = \sqrt{3} \cos x$
- $\tan x = \sqrt{3}$ *A1*
- $x = \frac{\pi}{3}, \frac{4\pi}{3}$ *A1*
- attempt to substitute **their** x into $f(x)$ *M1*
- e.g. $\cos\left(\frac{4\pi}{3}\right) + \sqrt{3} \sin\left(\frac{4\pi}{3}\right)$
- correct substitution *A1*
 e.g. $-\frac{1}{2} + \sqrt{3}\left(-\frac{\sqrt{3}}{2}\right)$
- correct working that clearly leads to -2 *A1*
 e.g. $-\frac{1}{2} - \frac{3}{2}$
- $q = -2$ *AG* *N0*
- (ii) correct calculations to find $f'(x)$ either side of $x = \frac{4\pi}{3}$ *A1A1*
- e.g. $f'(\pi) = 0 - \sqrt{3}, f'(2\pi) = 0 + \sqrt{3}$
- $f'(x)$ changes sign from negative to positive *R1*
- so A is a minimum *AG* *N0*
[10 marks]
- (c) max when $x = \frac{\pi}{3}$ *R1*
- correctly substituting $x = \frac{\pi}{3}$ into $f(x)$ *A1*
- e.g. $\frac{1}{2} + \sqrt{3}\left(\frac{\sqrt{3}}{2}\right)$
- max value is 2 *A1* *N1*
[3 marks]
- (d) $r = 2, a = \frac{\pi}{3}$ *A1A1* *N2*
[2 marks]

Total [17 marks]



MATHEMATICS
STANDARD LEVEL
PAPER 2

Candidate session number

0	0								
---	---	--	--	--	--	--	--	--	--

SPECIMEN

Examination code

1 hour 30 minutes

X	X	X	X	-	X	X	X	X
---	---	---	---	---	---	---	---	---

INSTRUCTIONS TO CANDIDATES

- Write your session number in the boxes above.
- Do not open this examination paper until instructed to do so.
- A graphic display calculator is required for this paper.
- Section A: answer all questions in the boxes provided.
- Section B: answer all questions on the answer sheets provided. Write your session number on each answer sheet, and attach them to this examination paper and your cover sheet using the tag provided.
- At the end of the examination, indicate the number of sheets used in the appropriate box on your cover sheet.
- Unless otherwise stated in the question, all numerical answers should be given exactly or correct to three significant figures.
- A clean copy of the **Mathematics SL formula booklet** is required for this paper.
- The maximum mark for this examination paper is [90 marks].




2. [Maximum mark: 7]

Let $f(x) = 4x - e^{x-2} - 3$, for $0 \leq x \leq 5$.

(a) Find the x -intercepts of the graph of f . [3 marks]

(b) On the grid below, sketch the graph of f . [3 marks]



(c) Write down the gradient of the graph of f at $x = 3$. [1 mark]

.....
.....
.....



5. [Maximum mark: 7]

The probability of obtaining heads on a biased coin is 0.4. The coin is tossed 600 times.

(a) (i) Write down the mean number of heads.

(ii) Find the standard deviation of the number of heads.

[4 marks]

(b) Find the probability that the number of heads obtained is less than one standard deviation away from the mean.

[3 marks]

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....



6. [Maximum mark: 7]

The following diagram shows a pole BT 1.6 m tall on the roof of a vertical building.
The angle of depression from T to a point A on the horizontal ground is 35° .
The angle of elevation of the top of the building from A is 30° .



Find the height of the building.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....



7. [Maximum mark: 8]

A circle centre O and radius r is shown below. The chord $[AB]$ divides the area of the circle into two parts. Angle AOB is θ .



- (a) Find an expression for the area of the shaded region. [3 marks]

- (b) The chord $[AB]$ divides the area of the circle in the ratio 1:7. Find the value of θ . [5 marks]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Do **NOT** write solutions on this page.

SECTION B (43 Marks)

Answer **all** the questions on the answer sheets provided. Please start each question on a new page.

8. [Maximum mark: 13]

Each day, a factory recorded the number (x) of boxes it produces and the total production cost (y) dollars. The results for nine days are shown in the following table.

x	26	44	65	43	50	31	68	46	57
y	400	582	784	625	699	448	870	537	724

(a) Write down the equation of the regression line of y on x . [2 marks]

Use your regression line as a model to answer the following.

(b) Interpret the meaning of

(i) the gradient;

(ii) the y -intercept. [2 marks]

(c) Estimate the cost of producing 60 boxes. [2 marks]

(d) The factory sells the boxes for \$19.99 each. Find the least number of boxes that the factory should produce in one day in order to make a profit. [3 marks]

(e) Comment on the appropriateness of using your model to

(i) estimate the cost of producing 5000 boxes;

(ii) estimate the number of boxes produced when the total production cost is \$540. [4 marks]



Do **NOT** write solutions on this page.

9. [Maximum mark: 16]

$$\text{Let } h(x) = \frac{2x-1}{x+1}, x \neq -1.$$

(a) Find $h^{-1}(x)$. [4 marks]

(b) (i) Sketch the graph of h for $-4 \leq x \leq 4$ and $-5 \leq y \leq 8$, including any asymptotes.

(ii) Write down the equations of the asymptotes.

(iii) Write down the x -intercept of the graph of h . [7 marks]

(c) Let R be the region in the first quadrant enclosed by the graph of h , the x -axis and the line $x = 3$.

(i) Find the area of R .

(ii) Write down an expression for the volume obtained when R is revolved through 360° about the x -axis. [5 marks]



Do **NOT** write solutions on this page.

10. [Maximum mark: 14]

A rock falls off the top of a cliff. Let h be its height above ground in metres, after t seconds.

The table below gives values of h and t .

t (seconds)	1	2	3	4	5
h (metres)	105	98	84	60	26

- (a) Jane thinks that the function $f(t) = -0.25t^3 - 2.32t^2 + 1.93t + 106$ is a suitable model for the data. Use Jane's model to
- (i) write down the height of the cliff;
 - (ii) find the height of the rock after 4.5 seconds;
 - (iii) find after how many seconds the height of the rock is 30 m. [5 marks]
- (b) Kevin thinks that the function $g(t) = -5.2t^2 + 9.5t + 100$ is a better model for the data. Use Kevin's model to find when the rock hits the ground. [3 marks]
- (c) (i) On graph paper, using a scale of 1 cm to 1 second, and 1 cm to 10 m, plot the data given in the table.
- (ii) By comparing the graphs of f and g with the plotted data, explain which function is a better model for the height of the falling rock. [6 marks]
-



Please **do not** write on this page.

Answers written on this page
will not be marked.



1212



MARKSCHEME

SPECIMEN

MATHEMATICS

Standard Level

Paper 2

Instructions to Examiners

Abbreviations

- M** Marks awarded for attempting to use a correct **Method**.
- A** Marks awarded for an **Answer** or for **Accuracy**; often dependent on preceding **M** marks.
- R** Marks awarded for clear **Reasoning**.
- N** Marks awarded for **correct** answers if **no** working shown.
- AG** Answer given in the question and so no marks are awarded.

Using the markscheme

1 General

Award marks using the annotations as noted in the markscheme eg **M1**, **A2**.

2 Method and Answer/Accuracy marks

- Do **not** automatically award full marks for a correct answer; all working **must** be checked, and marks awarded according to the markscheme.
- It is generally not possible to award **M0** followed by **A1**, as **A** mark(s) depend on the preceding **M** mark(s), if any. An exception to this rule is when work for **M1** is missing, as opposed to incorrect (see point 4).
- Where **M** and **A** marks are noted on the same line, e.g. **M1A1**, this usually means **M1** for an **attempt** to use an appropriate method (e.g. substitution into a formula) and **A1** for using the **correct** values.
- Where there are two or more **A** marks on the same line, they may be awarded independently; so if the first value is incorrect, but the next two are correct, award **A0A1A1**.
- Where the markscheme specifies **M2**, **N3**, etc., do **not** split the marks, unless there is a note.
- Once a correct answer to a question or part-question is seen, ignore further working.

3 N marks

If **no** working shown, award **N** marks for **correct** answers. In this case, ignore mark breakdown (**M**, **A**, **R**).

- Do **not** award a mixture of **N** and other marks.
- There may be fewer **N** marks available than the total of **M**, **A** and **R** marks; this is deliberate as it penalizes candidates for not following the instruction to show their working.
- If a candidate has incorrect working, which somehow results in a correct answer, do **not** award the **N** marks for this correct answer. However, if the candidate has indicated (usually by crossing out) that the working is to be ignored, award the **N** marks for the correct answer.

4 Follow through marks (only applied after an error is made)

Follow through (**FT**) marks are awarded where an incorrect answer from one **part** of a question is used correctly in **subsequent** part(s) or subpart(s). Usually, to award **FT** marks, **there must be working present** and not just a final answer based on an incorrect answer to a previous part. However, if the only marks awarded in a subpart are for the answer (i.e. there is no working expected), then **FT** marks should be awarded if appropriate.

- Within a question part, once an **error** is made, no further **A** marks can be awarded for work which uses the error, but **M** marks may be awarded if appropriate.
- If the question becomes much simpler because of an error then use discretion to award fewer **FT** marks.
- If the error leads to an inappropriate value (e.g. probability greater than 1, use of $r > 1$ for the sum of an infinite GP, $\sin \theta = 1.5$, non integer value where integer required), do not award the mark(s) for the final answer(s).
- The markscheme may use the word “their” in a description, to indicate that candidates may be using an incorrect value.
- Exceptions to this rule will be explicitly noted on the markscheme.
- If a candidate makes an error in one part, but gets the correct answer(s) to subsequent part(s), award marks as appropriate, unless the question says hence. It is often possible to use a different approach in subsequent parts that does not depend on the answer to previous parts.

5 Mis-read

If a candidate incorrectly copies information from the question, this is a mis-read (**MR**). Apply a **MR** penalty of 1 mark to that question

- If the question becomes much simpler because of the **MR**, then use discretion to award fewer marks.
- If the **MR** leads to an inappropriate value (e.g. probability greater than 1, use of $r > 1$ for the sum of an infinite GP, $\sin \theta = 1.5$, non integer value where integer required), do not award the mark(s) for the final answer(s).
- Miscopying of candidates’ own work does **not** constitute a misread, it is an error.

6 Alternative methods

Candidates will sometimes use methods other than those in the markscheme. Unless the question specifies a method, other correct methods should be marked in line with the markscheme

- Alternative methods for complete questions are indicated by **METHOD 1**, **METHOD 2**, etc.
- Alternative solutions for part-questions are indicated by **EITHER . . . OR**.

7 Alternative forms

Unless the question specifies otherwise, **accept** equivalent forms.

- As this is an international examination, accept all alternative forms of **notation**.
- In the markscheme, equivalent **numerical** and **algebraic** forms will generally be written in brackets immediately following the answer.
- In the markscheme, **simplified** answers, (which candidates often do not write in examinations), will generally appear in brackets. Marks should be awarded for either the form preceding the bracket or the form in brackets (if it is seen).

8 Accuracy of Answers

If the level of accuracy is specified in the question, a mark will be allocated for giving the answer to the required accuracy.

There are 2 types of accuracy errors, and the final answer mark should not be awarded if these errors occur.

- **Rounding errors:** only applies to final answers not to intermediate steps.
- **Level of accuracy:** when this is not specified in the question the general rule applies to final answers: unless otherwise stated in the question all numerical answers must be given exactly or correct to three significant figures.

9 Calculators

A GDC is required for paper 2, but calculators with symbolic manipulation features (e.g. TI-89) are not allowed.

Calculator notation

The Mathematics SL guide says:

Students must always use correct mathematical notation, not calculator notation.

Do **not** accept final answers written using calculator notation. However, do not penalize the use of calculator notation in the working.

10 Style

The markscheme aims to present answers using good communication, e.g. if the question asks to find the value of k , the markscheme will say $k = 3$, but the marks will be for the correct value 3 – there is *usually no need for the “ $k =$ ”*. *In these cases, it is also usually acceptable* to have another variable, as long as there is no ambiguity in the question, e.g. if the question asks to find the value of p and of q , then the student answer needs to be clear. Generally, the only situation where the full answer is required is in a question which asks for equations – *in this case the markscheme will say “must be an equation”*.

The markscheme often uses words to describe what the marks are for, followed by examples, using the e.g. notation. These examples are not exhaustive, and examiners should check what candidates have written, to see if they satisfy the description. Where these marks are **M** marks, the examples may include ones using poor notation, to indicate what is acceptable.

SECTION A

1. (a) attempt to substitute into sum formula for AP **M1**
e.g. $S_{20} = \frac{20}{2}(2(-7) + 19d)$, $\frac{20}{2}(-7 + u_{20})$
- setting up correct equation using sum formula **A1**
- e.g. $\frac{20}{2}(2(-7) + 19d) = 620$
 $d = 4$ **A1 N2**
[3 marks]
- (b) correct substitution $-7 + 77(4)$ **A1**
 $u_{78} = 301$ **A1 N2**
[2 marks]
- Total [5 marks]**

2. (a) intercepts when $f(x)=0$


M1

(0.827, 0) (4.78, 0) (accept $x=0.827$ $x=4.78$)

A1A1

N3

[3 marks]



A1A1A1

N3

Note: Award **A1** for maximum point in circle, **A1** for x-intercepts in circles, **A1** for correct shape (y approximately greater than -3.14).

[3 marks]

(c) gradient is 1.28


A1

N1

[1 mark]

Total [7 marks]

3. (a)



A1A1A1

N3

Note: Award **A1** for 450, **A1** for a to the right of the mean, **A1** for area 0.27.

[3 marks]

- (b) valid approach
 e.g. $P(X < a) = 1 - P(X > a)$, 0.73

 $a = 462.256\dots$

 $a = 462$

M1

A1

A1

N3

[3 marks]

Total [6 marks]

4. (a) L_1

A1

N1

[1 mark]

- (b) $\begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}$

A1

N1

[1 mark]

- (c) choosing correct direction vectors

A1A1

e.g. $\begin{pmatrix} 3 \\ -2 \\ 1 \end{pmatrix}, \begin{pmatrix} -1 \\ 2 \\ -a \end{pmatrix}$

recognizing that $\mathbf{a} \cdot \mathbf{b} = 0$

M1

correct substitution

A1

e.g. $-3 - 4 - a = 0$

$a = -7$

A1

N3

[5 marks]

Total [7 marks]

5.	(a)	(i) recognizing binomial with $n = 600$, $p = 0.4$	M1	
		$E(X) = 240$	A1	N2
		(ii) correct substitution into formula for variance or standard deviation	A1	
		e.g. 144 , $\sqrt{600 \times 0.4 \times 0.6}$		
		$sd = 12$	A1	N1
				[4 marks]
	(b)	attempt to find range of values	M1	
		e.g. 240 ± 12 $228 < X < 252$		
		evidence of correct approach	A1	
		e.g. $P(X \leq 251) - P(X \leq 228)$		
		$P(228 < X < 252) = 0.662$	A1	N2
				[3 marks]
				Total [7 marks]

6. METHOD 1

appropriate approach **M1**
e.g. completed diagram

attempt at set up **A1**
e.g. correct placement of one angle

$$\tan 30 = \frac{h}{x}, \tan 35 = \frac{h+1.6}{x} \quad \text{A1A1}$$

attempt to set up equation **M1**
e.g. isolate x

correct equation **A1**

$$\text{e.g. } \frac{h}{\tan 30} = \frac{h+1.6}{\tan 35}$$

$$h = 7.52 \quad \text{A1} \quad \text{N3} \quad \text{[7 marks]}$$

METHOD 2

$$\sin 30 = \frac{h}{1} \quad \text{A1}$$

in triangle ATB, $\hat{A} = 5^\circ$, $\hat{T} = 55^\circ$ **A1A1**
choosing sine rule **M1**

correct substitution

$$\text{e.g. } \frac{h / \sin 30}{\sin 55} = \frac{1.6}{\sin 5} \quad \text{A1}$$

$$h = \frac{1.6 \times \sin 30 \times \sin 55}{\sin 5} \quad \text{A1}$$

$$h = 7.52 \quad \text{A1} \quad \text{N3} \quad \text{[7 marks]}$$

7.	(a)	substitution into formula for area of triangle	A1
		e.g. $\frac{1}{2}r \times r \sin \theta$	
		evidence of subtraction	M1
		correct expression	A1 N2
		e.g. $\frac{1}{2}r^2\theta - \frac{1}{2}r^2 \sin \theta, \frac{1}{2}r^2(\theta - \sin \theta)$	
			[3 marks]
	(b)	evidence of recognizing that shaded area is $\frac{1}{8}$ of area of circle	M1
		e.g. $\frac{1}{8}$ seen anywhere	
		setting up correct equation	A1
		e.g. $\frac{1}{2}r^2(\theta - \sin \theta) = \frac{1}{8}\pi r^2$	
		eliminating 1 variable	M1
		e.g. $\frac{1}{2}(\theta - \sin \theta) = \frac{1}{8}\pi, \theta - \sin \theta = \frac{\pi}{4}$	
		attempt to solve	M1
		e.g. a sketch, writing $\sin x - x + \frac{\pi}{4} = 0$	
		$\theta = 1.77$ (do not accept degrees)	A1 N1
			[5 marks]
			Total [8 marks]

SECTION B

8.	(a)	$y = 10.7x + 121$	A1A1	N2	[2 marks]
	(b)	(i) additional cost per box (unit cost)	A1	N1	
		(ii) fixed costs	A1	N1	[2 marks]
	(c)	attempt to substitute into regression equation e.g. $y = 10.7 \times 60 + 121$, $y = 760.12\dots$ cost = \$760 (accept \$763 from 3 s.f. values)	M1		
			A1	N2	[2 marks]
	(d)	setting up inequality (accept equation) e.g. $19.99x > 10.7x + 121$ $x > 12.94\dots$ 13 boxes (accept 14 from $x > 13.02$, using 3 s.f. values)	M1		
			A1		
			A1	N2	
		Note: Exception to the FT rule: if working shown, award the final A1 for a correct integer solution for their value of x .			[3 marks]
	(e)	(i) this would be extrapolation, not appropriate	R1R1	N2	
		(ii) this regression line cannot predict x from y , not appropriate	R1R1	N2	[4 marks]
					Total [13 marks]

9. (a) $y = \frac{2x-1}{x+1}$

interchanging x and y (seen anywhere)

M1

e.g. $x = \frac{2y-1}{y+1}$

correct working

A1

e.g. $xy + x = 2y - 1$

collecting terms

A1

e.g. $x+1 = 2y - xy$, $x+1 = y(2-x)$


$h^{-1}(x) = \frac{x+1}{2-x}$

A1

N2

[4 marks]

(b) (i)



A1A1A1A1

N4

Note: Award **A1** for approximately correct intercepts,
A1 for correct shape, **A1** for asymptotes,
A1 for approximately correct domain and range.

(ii) $x = -1, y = 2$

A1A1

N2

(iii) $\frac{1}{2}$

A1

N1

[7 marks]

continued ...

Question 9 continued

(c) (i) area = 2.06

A2**N2**(ii) attempt to substitute into volume formula (do not accept $\pi \int_a^b y^2 dx$)**M1**

$$\text{volume} = \pi \int_{\frac{1}{2}}^3 \left(\frac{2x-1}{x+1} \right)^2 dx$$


A2**N3****[5 marks]****Total [16 marks]**

10.	(a)	(i)	106 m	A1	N1
		(ii)	substitute $t = 4.5$ $h = 44.9$ m	M1 A1	N2
		(iii)	set up suitable equation e.g. $f(t) = 30$ $t = 4.91$	M1 A1	N1 [5 marks]
	(b)		recognising that height is 0 set up suitable equation e.g. $g(t) = 0$ $t = 5.39$ secs	A1 M1 A1	N2 [3 marks]

continued ...

Question 10 continued

(c) (i)



A1A2

N3

Note: Award **A1** for correct scales on axes,
A2 for 5 correct points, **A1** for 3 or 4 correct points.

(ii) Jane's function, with **2** valid reasons
 e.g. Jane's passes very close to all the points, Kevin's has the rock clearly going up initially – not possible if rock falls

A1R1R1

N3

Note: Although Jane's also goes up initially, it only goes up very slightly, and so is the better model.

[6 marks]

Total [14 marks]